



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/561,029	03/16/2007	Douglas Spencer Millar	ALAR9.001APC	1871
20995 7590 04/13/2009 KNOBBE MARTENS OLSON & BEAR LLP 2040 MAIN STREET FOURTEENTH FLOOR IRVINE, CA 92614				
EXAMINER THOMAS, DAVID C				
ART UNIT 1637		PAPER NUMBER		
NOTIFICATION DATE 04/13/2009		DELIVERY MODE ELECTRONIC		

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Notice of the Office communication was sent electronically on above-indicated "Notification Date" to the following e-mail address(es):

jcartee@kmob.com
eOAPilot@kmob.com

Office Action Summary

Application No.

10/561,029

Applicant(s)

MILLAR, DOUGLAS SPENCER

Examiner

DAVID C. THOMAS

Art Unit

1637

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --
Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) OR THIRTY (30) DAYS, WHICHEVER IS LONGER, FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) ☒ Responsive to communication(s) filed on 05 March 2009.
2a) ☐ This action is **FINAL**. 2b) ☒ This action is non-final.
3) ☐ Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

Disposition of Claims

- 4) ☒ Claim(s) 1-29 is/are pending in the application.
4a) Of the above claim(s) 20-29 is/are withdrawn from consideration.
5) ☐ Claim(s) _____ is/are allowed.
6) ☒ Claim(s) 1-19 is/are rejected.
7) ☐ Claim(s) _____ is/are objected to.
8) ☐ Claim(s) _____ are subject to restriction and/or election requirement.

Application Papers

- 9) ☐ The specification is objected to by the Examiner.
10) ☐ The drawing(s) filed on _____ is/are: a) ☐ accepted or b) ☐ objected to by the Examiner.
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
11) ☐ The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

Priority under 35 U.S.C. § 119

- 12) ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
a) ☐ All b) ☐ Some * c) ☐ None of:
1. ☐ Certified copies of the priority documents have been received.
2. ☐ Certified copies of the priority documents have been received in Application No. _____.
3. ☐ Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

* See the attached detailed Office action for a list of the certified copies not received.

Attachment(s)

- 1) ☒ Notice of References Cited (PTO-892)
2) ☐ Notice of Draftsperson's Patent Drawing Review (PTO-948)
3) ☒ Information Disclosure Statement(s) (PTO-893)
Paper No(s)/Mail Date 6/30/2008, 1/5/2007
4) ☐ Interview Summary (PTO-413)
Paper No(s)/Mail Date _____
5) ☐ Notice of Informal Patent Application
6) ☐ Other: _____

DETAILED ACTION

Election/Restrictions

1. Applicant's election without traverse of Group I, claims 1-19, in the reply filed on March 5, 2009 is acknowledged. Claims 20-29 are withdrawn from further consideration pursuant to 37 CFR 1.142(b) as being drawn to a nonelected invention, there being no allowable generic or linking claim.

Claim Rejections - 35 USC § 103

2. The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

3. Claims 1, 3-5 and 8-13 are rejected under 35 U.S.C. 103(a) as being unpatentable over Dean et al. (Proc. Natl. Acad. Sci. U.S.A. (2002) 99:5261-5266, submitted on IDS) in view of Berlin, K. (U.S. Patent No. 7,008,770) and further in view of Olek (Nucleic Acids Res. 24:5064-5066, submitted on IDS).

With regard to claims 1, 3, 4, 8 and 9, Dean teaches a method for whole genome amplification (WGA) (for overview, see Abstract and p. 5261, column 1, second paragraph) comprising:

(b) providing a population of random X-mers of exonuclease-resistant primers capable of binding to at least one strand of the modified DNA, wherein X is an integer 3

or greater (exonuclease-resistant hexamers containing thiophosphate were provided, p. 5261, column 2, lines 10-11 and 33-34);

(c) contacting the modified DNA with nucleotides and a polymerase capable of amplifying double stranded DNA (a reaction mix was prepared containing human genomic DNA, dNTPs, hexamer primers and phi29 DNA polymerase, p. 5261, column 2, lines 30-37); and

(d) allowing the polymerase to amplify the modified DNA (multiple displacement amplification, MDA, occurred during incubation at 30°C for 18 hr, p. 5261, column 2, lines 37-38, p. 5263, column 1, lines 4-11 and Figure 1; the genomic DNA is not substantially degraded since there is no high temperature denaturation step used, p. 5263, column 2, lines 29-43).

With regard to claim 5, Dean teaches a method wherein the exonuclease-resistant primers are oligonucleotides or oligonucleotide analogues containing at least one intercalator pseudonucleotide forming an intercalating nucleic acid (INA) (hexamers are phosphorothioate-containing oligonucleotides, p. 5261, column 2, lines 10-11 and 33-34).

With regard to claims 10 and 11, Dean teaches a method wherein the primers contain from 3 to 40 bases, or about 6 to 20 bases (exonuclease-resistant hexamers containing thiophosphate were provided, p. 5261, column 2, lines 10-11 and 33-34).

With regard to claims 12 and 13, Dean teaches a method wherein the polymerase is selected from phi29, or a modified version thereof, or a functional equivalent thereof capable of amplifying double stranded DNA in vitro without the need

to denature the DNA (phi29 DNA polymerase is used in MDA reactions, p. 5261, column 2, lines 30-37, thus avoiding the need for denaturation of the genomic DNA template, p. 5263, column 2, lines 29-43).

Dean does not teach a method wherein genomic DNA is treated with a modifying agent which modifies cytosine bases but does not modify 5'-methyl-cytosine bases under conditions to form single stranded modified DNA, and wherein the agent is bisulphite, acetate or citrate, including sodium bisulphite. Dean does not teach a method wherein the primers are formed of two populations of INA primers comprising random X-mers containing either the bases A, G and T or the bases A, C and T, wherein one population of primers is capable of binding to one strand of DNA while the other population of primers is capable of binding to a complimentary synthesized strand of the DNA stand to which the first population of primers bind.

Berlin teaches a method for performing complex PCR amplifications such as whole genome amplifications (column 1, lines 7-12) wherein the template DNA is first treated with a reagent such as bisulphite that reacts differently with 5-methylcytosine and cytosine to change their base-pairing behavior (column 4, lines 37-44). Berlin also teaches a method wherein two populations of primers are used, one containing bases A, G and T and the other containing bases A, C and T, wherein one population binds to one strand while the other binds to the complementary strand and wherein at least one population can contain degenerate base positions (column 4, lines 33-35, column 5, lines 4-14 and column 6, lines 46-58).

Olek teaches an improved method for modification of DNA with sodium bisulphite for analysis of cytosine methylation in which the bisulphate treatment is performed on material embedded in agarose beads, resulting in formation of single-stranded modified DNA in the agarose since the denatured strands are prevented from renaturing in the agarose during chemical treatment (see Abstract and p. 5064, column 1, last line to column 2, line 5).

It would have been prima facie obvious to one having ordinary skill in the art at the time the invention was made to combine the whole genome amplification (WGA) methods of Dean and Berlin using nuclease-resistant hexamer primers to amplify genomic DNA by multiple displacement amplification (Dean) and amplifying bisulphite-treated DNA using primers containing 3 of 4 bases (Berlin) with the methods of Olek, who teaches sodium bisulphate treatment of DNA embedded in agarose to form and maintain the DNA in single-stranded form during treatment, since the methods of bisulphite treatment of Berlin and Olek can be easily adapted to the WGA methods of Dean with appropriately designed primers to amplify bisulphite-modified templates. Thus, an ordinary practitioner would have been motivated to combine the methods of Dean, Berlin and Olek since the hexamer primers taught by Dean can be designed to contain appropriate base compositions to amplify both the bisulphite-modified template and PCR products complementary to the original template. The bisulphite method of Berlin for performing WGA has the advantage by allowing identification of targets strands containing 5-methylcytosine versus cytosine bases by using primers which bind to the treated strand or the complement of the treated strand (Berlin, column 2, lines 39-

49, column 3, lines 37-43 and column 4, line 61 to column 5, line 14). Furthermore, the agarose methods of Olek increases sensitivity of detection by providing complete bisulphite treatment of the single-stranded template and also minimizes loss of material by eliminating DNA isolation or precipitation steps(Olek, p. 5064, column 1, last line to column 2, line 5 and p. 5065, column 2, line 12 to p. 5066, column 2, line 4). Moreover, Berlin indicates that one preferred method of bisulphite treatment for preparation of DNA for PCR and methylation analysis is the agarose method of Olek (Berlin, column 9, lines 14-26).

4. Claim 2 is rejected under 35 U.S.C. 103(a) as being unpatentable over Dean et al. (Proc. Natl. Acad. Sci. U.S.A. (2002) 99:5261-5266, submitted on IDS) in view of Berlin, K. (U.S. Patent No. 7,008,770) and further in view of Olek et al. (Nucleic Acids Res. 24:5064-5066, submitted on IDS) as applied to claims 1, 3-5 and 8-13 above, and further in view of Raizis et al. (Anal. Biochem. 226:161-166, submitted on IDS).

Dean, Berlin and Olek together teach the limitations of claims 1, 3-5 and 8-13, as discussed above. However, neither Dean, Berlin nor Olek teach a method wherein conditions used for treating genomic DNA with a cytosine base-modifying agent do not result in substantial DNA fragmentation.

Raizis teaches a method for bisulphite treatment of DNA that minimizes template degradation, including reduction of the time required to achieve completion of the bisulphite reaction using higher concentrations of bisulphite (p. 162, column 1, line 21 to column 2, line 34 and p. 165, column 2, line 15 to p. 166, column 1, line 5).

It would have been *prima facie* obvious to one having ordinary skill in the art at the time the invention was made to combine the whole genome amplification (WGA) methods of Dean, Berlin and Olek using nuclease-resistant hexamer primers to amplify genomic DNA by multiple displacement amplification (Dean) and amplifying bisulphite-treated DNA using primers containing 3 of 4 bases (Berlin) after bisulphite treatment by the methods of Olek with the bisulphite treatment by the methods of Raizis that minimizes template degradation since the methods of bisulphite treatment of Raizis can be easily adapted to the WGA methods of Dean, Berlin and Olek with appropriately designed primers to amplify non-degraded bisulphite-modified templates. Thus, an ordinary practitioner would have been motivated to combine the methods of Dean, Berlin and Olek with those of Raizis since the methods of Raizis avoid excessive depurination that causes fragmentation and reduction of full-length template molecules, and thus avoiding such fragmentation dramatically increases the sensitivity of detection of modified target sequences (Raizis, p. 161, column 2, lines 18-25 and p. 165, column 2, line 15 to p. 166, column 1, line 5). This method can be easily adopted to the bisulphite methods of Berlin and Olek for efficient preparation of samples for WGA.

5. Claims 6 and 7 are rejected under 35 U.S.C. 103(a) as being unpatentable over Dean et al. (Proc. Natl. Acad. Sci. U.S.A. (2002) 99:5261-5266, submitted on IDS) in view of Berlin, K. (U.S. Patent No. 7,008,770) and further in view of Olek (Nucleic Acids Res. 24:5064-5066, submitted on IDS) as applied to claims 1, 3-5 and 8-13 above, and further in view of Christensen et al. (U.S. Patent Pub. No. 2006/0014144).

Dean, Berlin and Olek together teach the limitations of claims 1, 3-5 and 8-13, as discussed above. However, neither Dean, Berlin nor Olek teach a method wherein the oligonucleotide or oligonucleotide analogues used as exonuclease-resistant primers are intercalating nucleic acids (INAs) and selected from the group consisting of subunits of DNA, RNA, peptide nucleic acid (PNA), hexitol nucleic acid (HNA), MNA, altritol nucleic acid (ANA), locked nucleic acid (LNA), cyclohexanyl nucleic acid (CAN), CeNA, TNA, (2'-NH)-TNA, nucleic acid based conjugates, (3'-NH)-TNA, α -L-Ribo-LNA, α -L-Xylo-LNA, β -D-Xylo-LNA, α -D-Ribo-LNA, [3.2.1]-LNA, Bicyclo-DNA, 6-Amino-Bicyclo-DNA, 5-epi-Bicyclo-DNA, α -Bicyclo-DNA, Tricyclo-DNA, Bicyclo[4.3.0]-DNA, Bicyclo[3.2.1]-DNA, Bicyclo[4.3.0]amide-DNA, β -D-Ribopyranosyl-NA, α -L-Lyxopyranosyl-NA, 2'-R-RNA, 2'-OR-RNA, α -L-RNA, and β -D-RNA.

Christensen teaches an oligonucleotide or oligonucleotide analogues comprising at least one intercalator pseudonucleotide (paragraph 24, lines 1-3) that increase nuclease stability of the oligonucleotide (paragraph 30), and include analogues such as PNA, HNA, MNA, ANA, LNA, TNA and others (paragraph 167, lines 1-23).

It would have been *prima facie* obvious to one having ordinary skill in the art at the time the invention was made to combine the whole genome amplification (WGA) methods of Dean, Berlin and Olek using nuclease-resistant hexamer primers to amplify genomic DNA by multiple displacement amplification (Dean) and amplifying bisulphite-treated DNA using primers containing 3 of 4 bases (Berlin) after bisulphite treatment by the methods of Olek with the teachings of Christensen for oligonucleotides containing intercalator analogues such as PNA, HNA, MNA, ANA, LNA, TNA, since these

analogues can be easily substituted in the exonuclease-resistant primers taught in the WGA methods of Dean using templates prepared by the bisulphite methods Berlin and Olek. Thus, an ordinary practitioner would have been motivated to combine the methods of Dean, Berlin and Olek with the teachings of Christensen since the intercalator analogues of Christensen (paragraph 167, lines 1-23), which also include phosphorothioates as also practiced by Dean for designing exonuclease-resistant primers (Dean, p. 5261, column 2, lines 10-11 and 33-34), not only can intercalate into a double helix at a predetermined position, but also increase the affinity of such oligonucleotides for DNA, substantially increase the specificity of hybridization, reduce cross- and self-hybridization and increase nuclease stability of the oligonucleotide (Christensen, paragraphs 25-30), all parameters that would be useful in whole genome amplification of a template nucleic acid.

6. Claims 14-19 are rejected under 35 U.S.C. 103(a) as being unpatentable over Dean et al. (Proc. Natl. Acad. Sci. U.S.A. (2002) 99:5261-5266, submitted on IDS) in view of Berlin, K. (U.S. Patent No. 7,008,770) and further in view of Olek (Nucleic Acids Res. 24:5064-5066, submitted on IDS) as applied to claims 1, 3-5 and 8-13 above, and further in view of Hogrefe et al. (U.S. Patent Pub. No. 2003/0143577).

Dean, Berlin and Olek together teach the limitations of claims 1, 3-5 and 8-13, as discussed above. However, neither Dean, Berlin nor Olek teach a method for WGA wherein the polymerase comprises a polymerase cocktail comprising a mixture of at least one proof-reading DNA polymerase selected from the group consisting of Pfu

polymerase, Pfu polymerase turbo, Vent polymerase, Vent exo- polymerase, Pwo polymerase, 9°NmDNA polymerase, Therminator, Pfx DNA polymerase, Expand DNA polymerase, rTth DNA polymerase, and DyNAzyme EXT Polymerase and at least one non proof-reading DNA polymerase selected from the group consisting of Taq polymerase, Taq polymerase Stoffel fragment, Advantage DNA polymerase, AmpliTaq, AmpliTaq Gold, Titanium Taq polymerase, KlenTaq DNA polymerase, Platinum Taq polymerase, and Accuprime Taq polymerase, wherein the ratio of proof-reading polymerase to non proof-reading polymerase is at least about 1:2-1:10 and wherein amplification is carried out by DNA thermal cycling.

Hogrefe teaches high fidelity DNA polymerase compositions comprising a wild-type non-proofreading DNA polymerase such as Taq DNA polymerase and a mutant DNA polymerase such as Pfu DNA polymerase with full 3' to 5' exonuclease activity but about 90% or more reduced DNA polymerization activity relative to wild-type activity (paragraph 12, lines 1-6 and paragraph 75, lines 1-7), wherein the polymerase composition can be used in PCR reactions requiring thermal cycling (paragraph 223, lines 1-4).

It would have been prima facie obvious to one having ordinary skill in the art at the time the invention was made to combine the whole genome amplification (WGA) methods of Dean, Berlin and Olek using nuclease-resistant hexamer primers to amplify genomic DNA by multiple displacement amplification (Dean) and amplifying bisulphite-treated DNA using primers containing 3 of 4 bases (Berlin) after bisulphite treatment by the methods of Olek with the teachings of Hogrefe for amplification of DNA using DNA

polymerase cocktails of both proof-reading and non-proof-reading polymerases since such cocktails provide both high polymerization activity and high levels of polymerization fidelity and are easily substituted in the WGA assays of Dean, Berlin and Olek. Thus, an ordinary practitioner would have been motivated to combine the methods of Dean, Berlin and Olek with the high fidelity DNA polymerase compositions taught by Hogrefe since Hogrefe provides a polymerizing system that achieves both high fidelity without compromising product yield needed for WGA since the proof-reading polymerase is modified to reduce polymerization and thus does not compete with the highly processive non-proof-reading enzyme (Hogrefe, paragraph 8, lines 1-11 and paragraph 66, lines 1-13). Hogrefe teaches that such high-fidelity polymerase blends are useful in a variety of applications requiring PCR (see paragraph 311), and thus would be highly suitable for applications such as WGA using modified primers and bisulphate-modified templates.

Summary

7. Claims 1-19 are rejected. No claims are allowable.

Correspondence

8. Any inquiry concerning this communication or earlier communications from the examiner should be directed to David C. Thomas whose telephone number is 571-272-3320 and whose fax number is 571-273-3320. The examiner can normally be reached on 5 days, 9-5:30.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Gary Benzion can be reached on 571-272-0782. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free).

/David C Thomas/
Examiner, Art Unit 1637

/Kenneth R Horlick/
Primary Examiner, Art Unit 1637